



Diurnal patterns and removal of selected elements in two Norwegian wastewater treatment plants with primary treatment

Farkas, Julia; Polesel, Fabio; Kjos, Marianne; Booth, Andy M.

Published in:
SETAC Europe 28th Annual Meeting - Abstract book

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Farkas, J., Polesel, F., Kjos, M., & Booth, A. M. (2017). Diurnal patterns and removal of selected elements in two Norwegian wastewater treatment plants with primary treatment. In *SETAC Europe 28th Annual Meeting - Abstract book* (pp. 351-351). [WE095] Society of Environmental Toxicology and Chemistry.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Diurnal patterns and removal of selected elements in two Norwegian wastewater treatment plants with primary treatment

Julia Farkas¹, Fabio Polesel², Marianne Kjos³, Andy M. Booth¹

¹SINTEF Ocean, Trondheim, Norway

²DTU Environment, Technical University of Denmark, Kgs. Lyngby, Denmark

³SINTEF Materials and Chemistry, Trondheim, Norway

E-mail contact: andy.booth@sintef.no

Keywords: elements, wastewater treatment, removal, diurnal variations, size fractions

Discharges from wastewater treatment plants (WWTPs) are a source for pollutants to the environment. Here we studied influent patterns and removal of selected elements in two full-scale WWTPs, Ladehamneren (LARA) and Høvringen (HØRA) in Trondheim, Norway. Both WWTPs have significant industrial loading contributions (up to 40% in LARA), employ preliminary and primary treatment steps, including chemically aided flocculation (CFeO₄S/polyamine in LARA, polyacrylamide in HØRA), and discharge directly into Trondheimsfjord. In a 7 day sampling campaign, 24 h composite samples of influent and effluent wastewater, as well as sludge samples, were taken to determine influent concentrations and removal of Al, P, S, Cr, Fe, Ni, Cu, Zn, As, Cd and Pd. To study release patterns and gain information about potential sources, diurnal variations of elemental concentrations were determined and modelled in 8 h composite samples of raw influent wastewater from morning, evening and night discharges.

Element concentrations in 24 h composite influent samples were highest for S>P and Al and lowest for Cd<As<Cr and Pd. Concentrations of Al, P, Cr, Cu and Cd were higher in HØRA than LARA, with Fe loadings being approximately double. Removal efficiencies varied between the analysed elements, and were highest for Al (86%), P (74%) and Cu (57%) in LARA, which utilises both inorganic and organic flocculants. In contrast, removal rates were below <50% for P, Cu and S in HØRA. However, in LARA, concentrations of Fe, Ni and S were significantly higher in the treated effluent compared to the raw influent, deriving from the use of inorganic flocculant. This was also reflected in Fe and S concentrations in treated sludge.

Elemental concentrations in 8 h composite samples mostly followed general diurnal discharge patterns, with higher concentrations in mornings and evenings and lower concentrations at night. In HØRA, concentrations of most elements further correlated well with total suspended solid concentrations (TSS), with the strongest correlations observed for P, S and Cu ($R^2 > 0.9$). Correlations with TSS were less pronounced in LARA, and were weakest for Pb, Fe and Cu ($R^2 < 0.6$), which can be potentially attributed to the higher industrial loading contributions in LARA. Enrichment factors were high for P>Cu>Zn>Cd>As, and were still above 10 for Cr and Ni in biosolids, indicating anthropogenic sources for these elements. Several elements also occurred as nano- and micron-sized particles.